



## PACKAGE LEAFLET: INFORMATION FOR THE USER

### STUD 100 Desensitizing Spray for Men Lidocaine 9.6% w.w

Read this entire leaflet carefully because it contains important information for you.

This medicine is available without prescription. However, you still need to use STUD 100 carefully to get the best results from it.

- Keep this leaflet. You may need to read it again.
- Ask your pharmacist if you need more information or advice.
- If any of the side effects become serious, or if you notice any side effect not listed in this leaflet, please tell your doctor or pharmacist.

#### In this leaflet:

1.	What STUD 100 is and what it is used for
2.	Before you use STUD 100
3.	How to use STUD 100
4.	Possible side effects
5.	How to store STUD 100
6.	Further information

#### 1. WHAT STUD 100 IS AND WHAT IT IS USED FOR

The active ingredient, Lidocaine, is a local anaesthetic which is used to reduce the sensitivity to touch of the penis just before intercourse. The reduction in sensitivity helps to delay ejaculation (climax) in cases of over-rapid or premature ejaculation (coming to a climax too quickly).

#### 2. BEFORE YOU USE STUD 100

##### **X** Do not use STUD 100:-

- If you or your partner are allergic (hypersensitive) to Lidocaine or any of the other ingredients of STUD 100.
- If you or your partner are allergic (hypersensitive) to other local anaesthetics.
- If your partner is pregnant
- Do not use on broken or inflamed skin
- Do not spray near the eyes or nostrils

##### **i** Take special care with STUD 100

###### Elderly

STUD 100 is not recommended for use by the elderly

###### Kidney and Liver Problems

If you have, or have ever had, liver or kidney problems – ask your doctor before using the spray

##### **i** Using other medicines

High blood levels of the active ingredient of STUD 100, Lidocaine, can cause interactions with other medicines. Although this is unlikely to occur with the low levels absorbed from STUD 100 you should ask your doctor or pharmacist for advice before using the spray if you are taking other medicines.

Please tell your doctor or pharmacist if you are taking or have recently taken any other medicines, including medicines or herbal therapies obtained without a prescription.

##### **X** Pregnancy and breast-feeding

Although STUD 100 is applied to the penis only, it must not be used if your partner is pregnant.

Ask your doctor or pharmacist for advice before using any medicine.

###### **Driving and using machines**

STUD 100 is unlikely to impair the ability to drive or operate machinery

4092B

### 3. HOW TO USE STUD 100

The quantity used and timing will vary with individual requirements but the usual recommended dosage is 3 to 8 sprays to the head and the shaft of the penis 5 to 15 minutes before intercourse. You should always use the minimum effective quantity.

- This spray is for external use only. Do not spray near the eyes or nostrils.
- Wash off the spray after intercourse.
- Do not use the spray for more than 3 months without consulting your doctor.
- Do not exceed a maximum of 24 sprays in 24 hours.



#### **i** If you use more STUD 100 than is recommended

In the unlikely event of a suspected overdose seek prompt medical advice.

If you have any further questions on the use of this product, please ask your doctor or pharmacist.

### 4. POSSIBLE SIDE EFFECTS

STUD 100 is suitable for most men, but like all medicines it can sometimes cause side effects.

- Rash or irritation - If you or your partner develops a rash or irritation stop using the spray.

**i** No other side effects associated with the use of STUD 100 Desensitizing Spray for Men have been reported as only extremely low levels of Lidocaine are absorbed. However, high blood levels of Lidocaine (i.e. 25 times the recommended dose of STUD 100) can cause side effects such as nervousness, dizziness, convulsions, loss of consciousness, as well as effects on the heart and lungs.

If any of the side effects become serious, or if you notice any side effects not listed in this leaflet, please tell your doctor or pharmacist.

### 5. HOW TO STORE STUD 100

Keep out of the reach and sight of children.

- Do not use STUD 100 after the expiry date which is stated on can.
- The expiry date refers to the last day of that month.
- Store below 25°

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

### 6. FURTHER INFORMATION

#### **What STUD 100 contains:**

- The active substance is Lidocaine Base USP 9.6% w/w
- The other ingredients are Isopropyl Myristate, Stearic Acid, 2-(2-Ethoxyethoxy)ethanol and fragrance

#### **What STUD 100 looks like and contents of the pack**

Aluminium can with metered pump spray and aluminium cap, containing 12g STUD 100 Desensitizing Spray for Men.

#### **Marketing Authorisation Holder and Manufacturer**

Pound International Ltd  
47b Welbeck Street  
London  
W1G 9XA  
England

This product is manufactured on behalf of Pound International Ltd, by Fleet Laboratories Limited, 94 Rickmansworth Road, Watford, Herts. WD18 7JJ, United Kingdom

#### **This medicinal product is authorised in the Member States of the EEA under the following names:**

United Kingdom – STUD 100 Desensitizing Spray for Men

**This leaflet was last approved in 05/2017**

4092B

SO48498\_4092B\_P2.indd 09/05/2017 07:37